

EDTA Iron 5%

For Prevention or Correction of Iron
Deficiency in Most Crops

GUARANTEED ANALYSIS

Iron (Fe).....5.00%
5.0% Chelated Iron

Derived From: Iron, Ethylenediaminetetraacetate (Fe EDTA)

Information about the components of this fertilizer material may be obtained by writing to Cromartie Agricultural Chemicals, Inc. P.O. Box 71647, Albany, GA 31708, and giving the lot number which is found on the container.

Net Weight:

Pail: 50lb (29.49 kg)

Jug: 25lbs (12.24 kg)

Cs 2x2-1/2: 50lb (24.49 kg)

Net Contents:

Pail: 5 gals (18.92 liters)

Jug: 2-1/2 gals (9.46 liters)

Cs 2x2-1/2: 5 gals (18.92 liters)

Bulk _____ Lot Number: _____

GENERAL INFORMATION

EDTA Iron 5% contains fully chelated iron in a form readily available to plants. It can be effectively used as an additive to dry or liquid fertilizer in irrigation systems or as a foliar spray. **EDTA Iron 5%** is also compatible with most herbicides, insecticides and agricultural chemicals that may be utilized in combination with liquid fertilizer. **EDTA Iron 5%** is NOT compatible with CAN-17 or N-PHURIC ®

EDTA Dual Crop is most effective when placed in the root zone during planting or moved into the soil by irrigation, tillage or rainfall. It is important that the iron deficiency be treated before visual symptoms appear to ensure proper yields. Rates of application will vary depending upon the severity of the deficiency, climate and soil conditions and method of application. Consult your State Agricultural Experiment Station or Extension specialists for advice in selecting treatments from this label to best fit local conditions.

CAUTION: If label instructions are not followed for foliar applications, phytotoxicity may occur. If there is any doubt, apply the spray solution to a small test area of the crop or foliage to assess any undesirable effects or phytotoxicity before general application.

CAUTION: Harmful if swallowed. Avoid contact with eyes, skin, and clothing. Avoid breathing spray mist.

READ LABEL – FOLLOW DIRECTIONS!

Refer to MSDS for health, safety and environmental information.

KEEP OUT OF REACH OF CHILDREN

CAC

**CROMARTIE
AGRICULTURAL
CHEMICALS, INC.**

1319 US Hwy 82 West • Albany, GA 31708

DIRECTIONS FOR USE

Soil Application: **EDTA Iron 5%** can be applied alone or in combination with dry or liquid fertilizer for pre-plant, starter or side dress applications. For most effective treatment of row crops, make soil applications in a band at planting time or as a side dressing shortly after planting.

Foliar Application: **EDTA Iron 5%** when diluted with water according to label recommendations, can be applied as a foliar spray. To ensure even distribution on leaf surface and increase effectiveness of foliar application, a small amount of wetting agent or detergent may be added. Aerial applications can be made at a concentration of 1 qt. per 5 - 10 gallons of water.

For Foliar or Soil Application

Recommendations for Use

Field and vegetable crops including corn, rice, potatoes, beans, cotton, tomatoes, etc.

Soil Application: Apply 2 - 4 quarts of **EDTA Iron 5%** per acre in liquid starter fertilizer. Rate will depend on extent of the iron deficiency and soil conditions.

Foliar Application: Apply 1 quart of **EDTA Iron 5%** in 20-100 gallons of water per acre to thoroughly wet foliage. Apply 4-6 weeks after planting or when iron deficiencies first appear.

Fruit and nut crops including almonds, walnuts, deciduous fruits, citrus and grapes.

Soil Application: Apply 4 quarts of **EDTA Iron 5%** per acre depending on soil conditions, size of tree and severity of deficiency.

Foliar Application: Apply 1 quart of **EDTA Iron 5%** per 20 to 100 gallons of water per acre and spray to thoroughly cover foliage 10 days to 2 weeks prior to bloom. Follow with a 2nd application in two weeks if required.

WHAT TO DO IN CASE OF CONTACT

If Ingested: Give large amounts of water or milk, if conscious. Contact a physician immediately.

Eye Contact: Flush eyes with clean water for 15 minutes. Seek medical attention immediately.

Skin Contact: Remove contaminated clothing and wash skin with soap and water. Seek medical attention if irritation develops or persists.

If Inhaled: Move to fresh air and seek medical attention if irritation develops or persists.

If there is contact made with spray solution containing pesticides and/or spray adjuvants, follow the "statement of Practical Treatment" on the product labeling.

This product may be corrosive to aluminum, mild steel and brass. Store in HDPE, fiberglass or stainless steel containers. Use only stainless steel, P.V.C., or polypropylene fittings.

Storage: Store in cool place in original container in a locked storage area inaccessible to children, pets, domestic animals or wildlife.

Container Disposal: Do not reuse empty containers. Triple rinse (or equivalent) then offer for recycling or disposal by procedures approved by state or local authorities.

CONDITIONS OF SALE: Cromartie Ag. Chemicals, Inc. (Company) warrants that this product conforms to the chemical description of the label and is reasonably fit for the purposes stated on the label when used in accordance with the directions under normal use. This warranty does not extend to the use of this product contrary to label instruction, or under abnormal use conditions, or under conditions not reasonably foreseen by the Company. THE COMPANY DISCLAIMS ALL OTHER WARRANTIES, EXPRESSED OR IMPLIED, INCLUDING ANY WARRANTY OF FITNESS OR MERCHANTABILITY. THE COMPANY SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT AND THE COMPANY'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE. BUYER AND SELLER ACKNOWLEDGE AND ASSUME ALL RISKS AND LIABILITY RESULTING FROM HANDLING, STORAGE AND USE OF THIS PRODUCT. THE COMPANY DOES NOT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTY, GUARANTEE OR REPRESENTATION CONCERNING THIS PRODUCT.